

VIRUS, Previous Infection

#532

Cf. #362
Activated: 5/1/63
Renewed: 5/1/64
Renewed: 5/1/65

THE COUNCIL FOR TOBACCO RESEARCH - U.S.A.

SUCCESSOR TO THE
TOBACCO INDUSTRY RESEARCH COMMITTEE

COMMITTEE:

Dr. Jacobson, Chm.
Dr. Lynch
Dr. Little

633 THIRD AVENUE
NEW YORK, N.Y. 10017

Application For Research Grant

Date: January 31, 1966

1. Name of Investigator: JOHN P. MANOS, M.D.

Associate in Microbiology

2. Title:

MEDICAL COLLEGE OF SOUTH CAROLINA.

3. Institution &

Address:

80 Barre Street
Charleston, South Carolina 29401

4. Project or Subject: THE ANTIGENIC STUDY OF THE VARIOUS NEOPLASTIC SUBCELLULAR PARTICLES
FOLLOWING SEPARATION BY ULTRA-FILTRATION METHODS.

5. Detailed Plan of Procedure (Use additional pages if more space is required.)

That neoplastic cells have an antigenic makeup distinct at least in part from normal cells has been a source of much interest and investigation. The significance of such studies for potential diagnostic, therapeutic, preventive and etiologic evaluations is obvious. Some pertinent references are listed.

As an outgrowth of a project currently in progress, it was felt that an antigenic study of subcellular fractions of neoplastic cells by methods used in the other project, i.e. differential ultra-filtration, the emergence of specific 'cancer' antigens might be effected. As indicated in progress report #4, work has been begun in this area. Cells from a human lung cancer and normal human lung have been disrupted mechanically by grinding and the subcellular material fractionated by ultra-filtration methods using the Gelman filterfuge apparatus.

These fractions, 10 from the human lung carcinoma and 10 from normal human lung, have been inoculated in paired rabbits using Freund's complete adjuvant and an intensive immunization schedule. The rabbits were exsanguinated and the cell fractions are being tested for antigenicity. The first method of approach is that of gel double immunodiffusion. Preliminary work by this method has shown that some of the fractions so far tested, both from cancer and normal lung cell are antigenic. It is planned, by cross absorption techniques to determine if any of the cancer cell fractions have or lack antigenic component(s) absent or present in the normal cell fractions. Other methods to be used for determining antigen specificity are the tanned red cell and possibly the complement-fixation techniques. Also, other approaches to the preparation of fractions could be used, such as density gradient centrifugation and column chromatography.

1005101310

BIBLIOGRAPHY

Bjorklund, B., et al., Antigenicity of pooled malignant and normal tissues by cytoimmunological technique. III. Distribution of tumor antigen. *J. Natl. Cancer Inst.* 26:433-45, 1961

Burnet, M., Innumological factors in the process of carcinogenesis. *Brit. Med. Bull.*, 20:154, May, 1964

Gorer, P. A., The Antigenic structure of Tumors. *Advances in Immunology*, Academic Press, New York, 1961, pp. 345-393

Habel, K., Antigenic Properties of Cells Transformed by Polyoma Virus. *Cold Spring Harbor Symposia on Quantitative Biology*, 27:433-439, 1962

Habel, K., Immunologic Aspects of Oncogenesis by Polyoma Virus, *Symposium on Fundamental Cancer Research*, Harper and Row, New York, 1963, p. 486-502

Old, L. J., et al., Antigenicity of Chemically Induced Tumors. *Ann. N. Y. Acad. Sci.* 101(1):80, 1962

Old, L. J., et al., Immunology of Experimental Tumors. *An. Rev. Med.*, 15:167, 1964

Prehn, R. T., Isoantigenicities of Chemically Induced Tumors. *Ann. N. Y. Acad. Sci.* 101(1):107, 1962

Prehn, R. T., Role of Immunologic Mechanisms in the Biology of Chemically and Physically Induced Tumors. *Symposium on Fundamental Cancer Research*, Harper and Row, New York, 1963, pp. 475-485

Sjogren, H. O., Experimental Immunization against Carcinogen-Induced and Virus-Induced Mouse Tumors in Isologous or Autologous Systems. *Symposium on Fundamental Cancer Research*, Harper and Row, New York, 1963, pp. 459-474

Zibler, L. A., Study of the Tumor Specificity of Gastric Cancer Tissue. *Ann. N. Y. Acad. Sci.*, 101(1):264, 1962

1005101311

6. Budget Plan:

a. Salaries	5000.00
b. Expendable Supplies	1000.00
c. Other Expenses	3000.00
d. Permanent Equipment	900.00
e. Overhead (15% of a, b, c)	990.00
Total	

7. Anticipated Duration of Work:

3 Years

8. Facilities and Staff Available:

Millipore filter apparatus and filters, Gelman centrifuge filter holders and filter, various centrifuges, including the Serval Regrigerated super-speed centrifuge, spectrophotometers, Micro Kjeldahl equipment, photographic equipment, chromatographic apparatus, sonic oscillators, fluorescent microscopy equipment, electron microscopy and other equipment pertinent to microbiological work. Apparatus for double immunodiffusion techniques and other serological techniques. Column chromatographic apparatus.

9. Additional Requirements: Need for a preparative ultra-centrifuge in the future.

10. Additional Information (including relation of work to other projects and other sources of support):

The methods used for subcellular fraction preparation derived from current project studying similar type fractions from mouse lung adenomas in an attempt to demonstrate viral activity in mice and tissue cultures. This latter project is presently being supported by The Council For Tobacco Research-U.S.A.

Signature

John J. Manos, M.D.
Director of Project

Business Officer of the Institution

1005101312